

## Kerry Basic

Benefit :

1. Improved soil structure.
2. Strong root formation, resistance to drought.
3. Increased yield.
4. Eliminates trace element deficiency.
5. Resistance to disease.
6. For ornamentals - Stronger colour, longer lasting blooms.

## 佳丽加基素

益处：


1. 改良土壤构造。
2. 加强根部构造，抵抗干旱季节。
3. 增加产量。
4. 释放微量元素的不足。
5. 抵抗病害侵袭。
6. 使观赏植物颜色鲜艳，开花更持久。

Net Content : **1 liter**

Manufacturing Date :

# Kerry Basic

佳麗  
基素



## Natural Seaweed Extract 纯海藻有机养素

Seaweed	10% w/v
Protein	1% w/v
Carbohydrate	10% w/v
40 Trace Elements	Al, Co, I, Mo, Na, Ni, Si, etc
Growth Stimulants	Betaines, Auxin, Cytokinins, Gibberelin

Imported & Distributed by :



**Mycrop Sdn Bhd** (507146-W)

36, Lalan Perindustrian Silibin 2, Off Jalan Jelapang, 30020 Ipoh,  
Perak Darul Ridzuan, Malaysia.  
Tel: (6) 05-5285281 Fax: (6) 05-5284281 Website : www.mycrop.com.my  
E-mail : mycrop@po.jaring.my

Foliar Application (ml/4.5 liter) 施用方法(毫升/4.5公升)

CROP 作物	TIMING			
	First - 5-10 ml/ 毫升	Second 第二 5-10 ml/ 毫升	Third 第三 7.5-10 ml/ 毫升	Optional 施用次数 7.5 ml/ 毫升
<b>Vegetables</b> 蔬菜 Cabbages, Cauliflower Brassica, Onion, Lettuce. 包菜, 花椰菜, 十字花科, 落, 生菜	At 4-6 leaf Stage. 4-6 月一时	10 days later. 10 天簽	At head Initiation 由头灌溉	-
<b>Fruit Vegetables</b> 果菜 Tomato, Peppers. 番茄, 甜椒	At 15-20cm growth 生长至15-20	At first bloom 開花初期	At first fruit set. 给果初期	Every 14 days
Beans, Peas. 豆类, 豌豆	At 4-6 leaf stage. 4-6 片叶时	At first bloom 開花初期	At first pods. 给豆荚初期	-
Watermelon, Muskmelon 西瓜, 蜜瓜	At 15-20cm growth. 生长至15-20 公分时	At first bloom 開花初期	At first fruit set. 给果初期	Within 48 hours of harvest. 选婚后48 小时内
Cucumber 黄瓜	At 4 leaf stage. 4片叶时	At first bloom 開花初期	7-14 days later 7-14 天后	-
<b>Fruits Tree</b> 果树 Mango, Pomelo, Guava etc.. 芒果, 拍子, 番 石榴等等..	Flowering bud formation, 花蕾形成时	Flowering 开花时	After 14 days. 14 天后	-
<b>Paddy</b> 稻谷	2 weeks after sowing 播种後2星期	Flowering 开花时期	After 14 days. 14 天后	-