

## Yield-CaB

Contains key nutrients of Calcium, Magnesium, Boron, Nitrogen and Zinc.

- a) Stimulates active and strong flower pollens.
- b) Enhances fruit setting.
- c) Enhances development and growing of cells.
- d) Builds strong cell wall, increases resistance to fruit rot and fruit cracking.
- e) Improves fruit quality and shelf life.

Recommendation for application :

Crop	Rate	Application
<b>Tropical fruits</b> Mango, Watermelon, Guava, Waxapple	5-7.5 ml/ 4.5 liter	- Apply 1-2 weeks before flowering. - Apply during blooming and early fruit setting.
<b>fruit vegetables</b> Tomato, Chili, Okra, Cucumber, Beans.		- Apply every 7-10 days during fruiting period.
<b>Leafy vegetables</b> Brassica sp., Kailan, Amaranth sp.		- Apply 7-10 days before harvesting.
Rice	1 L/ha	- Apply at panicle Initial & Milking Stage

# Yield-CaB

優



## Plant Nutrients 植物营养素

CaO 鈣 .....	10.5 %	MgO 鎂 .....	2.5 %
B <sub>2</sub> O <sub>5</sub> 硼 .....	2.5 %	N 氮 .....	1.5 %
Zn 鋅 .....		1.5 %	

Net Content : **1 liter**      Manufacturing Date :

Sole Distributor : **Mycrop Sdn Bhd (507146-W)**

36, Laluan Perindustrian Silibin 2, Off Jalan Jelapang, 30020 Ipoh, Perak Darul Ridzuan, Malaysia.

Tel: (6) 05-5285281 Fax: (6) 05-5284281 Website : [www.mycrop.com.my](http://www.mycrop.com.my)  
E-mail : [mycrop@po.jaring.my](mailto:mycrop@po.jaring.my)



優

含有微量元素如鈣, 鎂, 硼, 氮及鋅。

- a) 促進花粉活性及提高著果率。
- b) 提高著果量。
- c) 促進細胞分化及生長。
- d) 增強細胞壁, 提高抗病蟲害能力如腐果病及裂。
- e) 提高果實品質及增強儲藏能力。

推薦使用方法:

作物	用量	施用時期
熱帶水果 芒果, 西瓜, 番石榴, 蓮霧	5-7.5 毫升/ 4.5公升	- 開花前1-2星期使用. - 開花結果時期.
果菜 番茄, 辣椒, 羊角豆, 黃瓜, 豆類		結果期間每隔7-10天 使用
葉類 十字花科, 芥藍, 莧菜		采收前7-10天使用
稻米	1公升/ 公頃	幼穗形成期及孕穗期.